



Chelan County Population = 67,000 (29% under age 18)

Adult Tobacco Use**Current Cigarette Smoking (1987-1998)****Current Smokeless Tobacco Use (1993-1997)**

Chelan County
Washington State

Entire adult population
2.0% (95%CI: .6, 6.7)
2.9% (95%CI: 2.5, 3.8)


Males only
3.6% (95%CI: 1.1, 12.0)
5.7% (95%CI: 5.0, 6.5)

Youth Tobacco Use (1998)**Summary of the problem**

- ☐ Adult tobacco use is very similar to adult tobacco use throughout the state.
- ☐ Youth tobacco use for the Eastern Washington region is elevated above the state rate for both cigarettes and smokeless tobacco, among all age groups. More data collection is needed to describe Chelan County youth specifically.
- ☐ Maternal smoking during pregnancy is consistently and significantly lower than the state as a whole, and exhibits the same declining trend seen throughout the state. Chelan County is among the lowest-risk counties in Washington for maternal smoking.
- ☐ Retailer compliance is significantly worse than the state as a whole. Chelan County is among the highest-risk counties in Washington for retailers selling tobacco to minors.

Estimated number of current adult and youth smokers who will die prematurely as a result of tobacco use: 4,300


Smoking During Pregnancy (1987-1998)



Infants born annually to mothers who smoke during pregnancy: 43

Minors Access to Tobacco (1998-1999)

Percent Sales to Minors



Number of tobacco retailers in county: 116

Health at Risk (Estimated numbers for county)

Adults who currently smoke	11,800
Adults who currently use smokeless tobacco	1,400
Youth who currently smoke	1,400
Youth who currently use smokeless tobacco	600
ESTIMATED ANNUAL DEATHS DUE TO TOBACCO USE	110

Current Tobacco Prevention and Control Program Activities

Highlights during 1999 for Chelan and Douglas Counties

- ✓ Trained 120 high school youth as peer educators. Teens gave 34 presentations to 4th, 5th, and 6th grade students.
- ✓ Provided tobacco cessation program to middle school, Juvenile Detention, and PIC/Pentad (community program) youth.
- ✓ Partnered with cities of Cashmere, Leavenworth, and Wenatchee to pass Tobacco Ordinance Laws.

Annual Funding for Program (12-month funding amounts, FY 1999)

Source	Amount
Youth Access (Retailer License Fees)	\$ 9,119
CDC Grant	\$ 4,653
Total Per Capita Funding	\$.21


Agency receiving funding

Chelan-Douglas TOGETHER!
Chelan-Douglas TOGETHER!
CDC Recommended Community Per Capita Funding: \$.70-2.00

Local Contacts

Renee Hunter, Executive Director
Chelan-Douglas TOGETHER!
P.O. Box 3525
Wenatchee, WA 98807
(509) 662-7201
fax: (509) 664-7205
email: together@televar.com

Pat Malone, Administrator
Chelan-Douglas Health District
P.O. Box 429
Wenatchee, WA 98807-0429
(509) 886-6480
fax: (509) 886-6478
malonep@health-chelan-douglas.wa.gov



TECHNICAL NOTES: (ADDITIONAL DOCUMENTATION IN APPENDIX B)

TOTAL DEATHS DUE TO TOBACCO USE ARE ESTIMATED AS 19% OF ALL DEATHS (MCGINNIS AND FOEGE 1993); "DUMBBELL" BARS AROUND DATA POINTS FOR SURVEY DATA REPRESENT 95% CONFIDENCE INTERVALS – THE TRUE VALUE MAY BE ANYWHERE WITHIN THE BARS, BUT MAY NOT BE EXACTLY AT THE DATA POINT; NUMBER OF ADULTS/YOUTH WHO WILL DIE PREMATURELY FROM TOBACCO USE IS ESTIMATED AS 33% OF CURRENT ADULT SMOKERS (MMWR 1996), AND 21% OF CURRENT YOUTH SMOKERS –WHO MAY 'GROW OUT OF' SMOKING BEHAVIOR (USDHHS 1994); "CURRENT ADULT/YOUTH TOBACCO USERS" ARE SYNTHETIC ESTIMATES BASED ON STATEWIDE AGE-SPECIFIC PREVALENCE; FUNDING LEVELS FOR COUNTIES WERE CALCULATED BASED ON DEDICATED YOUTH TOBACCO PREVENTION FUNDS AND CDC FUNDS MANAGED BY THE DEPARTMENT OF HEALTH – COUNTIES MAY SUCCESSFULLY SEEK ADDITIONAL FUNDING FOR TOBACCO CONTROL FROM OTHER SOURCES; "HIGHLIGHTS DURING 1999" WERE IDENTIFIED BY LOCAL PROGRAM STAFF; "CDC RECOMMENDED FUNDING" IS BASED ON *BEST PRACTICES FOR TOBACCO CONTROL PROGRAMS* (CDC 1999).